

AD-A153 887 ON MGR (UPSILON) MULTIGRID METHODS(U) WISCONSIN

1/1


D KAMOWITZ ET AL. JAN 85 CSTR-575 AFOSR-TR-85-0351

AFOSR-82-0275

F/G 12/1

NL

END



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A


②

# COMPUTER SCIENCES DEPARTMENT

University of Wisconsin-  
Madison

AD-A153 887

DTIC  
ELECTE  
MAY 17 1985  
S B



ON MGR[v] MULTIGRID METHODS

by

David Kamowitz  
and  
Seymour V. Parter

Computer Sciences Technical Report #575

January 1985

DTIC FILE COPY

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION <b>UNCLASSIFIED</b>			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited.		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report #575			5. MONITORING ORGANIZATION REPORT NUMBER(S) <b>AFOSR-TR- 85 - 0351</b>		
6a. NAME OF PERFORMING ORGANIZATION University of Wisconsin		6b. OFFICE SYMBOL (If applicable)		7a. NAME OF MONITORING ORGANIZATION Air Force Office of Scientific Research	
6c. ADDRESS (City, State and ZIP Code) Computer Sciences Department Madison WI 53706			7b. ADDRESS (City, State and ZIP Code) Directorate of Mathematical & Information Sciences, Bolling AFB DC 20332-6448		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR		8b. OFFICE SYMBOL (If applicable) NM		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR-82-0275	
8c. ADDRESS (City, State and ZIP Code) Bolling AFB DC 20332-6448			10. SOURCE OF FUNDING NOS.		
			PROGRAM ELEMENT NO. 61102F	PROJECT NO. 2304	TASK NO. A3
11. TITLE (Include Security Classification) ON MGR[v] MULTIGRID METHODS					
12. PERSONAL AUTHOR(S) David Kamowitz and Seymour V. Parter					
13a. TYPE OF REPORT Technical		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Yr., Mo., Day) JAN 85	
15. PAGE COUNT 34					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB. GR.			
19. ABSTRACT The MGR[v] algorithm of Ries, Trottenberg and Winter with $v = 0$ and the Algorithm 2.1 of Braess are essentially the same multigrid algorithm for the discrete Poisson equation: $-\Delta_h U = f$ . In this report we consider the extension to the general diffusion equation $-\nabla \cdot p \nabla u = f$ , $p = p(x,y) \geq p_0 > 0$ . In particular, for the two-grid scheme we reobtain the basic result $\rho \leq \frac{1}{2}(1+Kh)$ in the stronger form $\ MG\ _{L_h} \leq \frac{1}{2}(1+Kh)$ . Computational results indicate that other constant coefficient results carry over as well.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL CPT John P. Thomas, Jr.			22b. TELEPHONE NUMBER (Include Area Code) (202) 767- 5026		22c. OFFICE SYMBOL NM

ON MGR[v] MULTIGRID METHODS\*

by

David Kamowitz  
and  
Seymour V. Parter

DTIC  
ELECTE  
MAY 17 1985  
S B D

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH  
MAIL ROOM  
Chief, Technical Support Division

\*Supported by the Air Force Office of Scientific Research under contract  
#AFOSR-82-0275 and the Los Alamos National Laboratory.

# ABSTRACT

The MGR[v] algorithm of Ries, Trottenberg and Winter with  $v = 0$  and the Algorithm 2.1 of Braess are essentially the same multigrid algorithm for the discrete Poisson equation:  $-\Delta_h U = f$ . In this report we consider the extension to the general diffusion equation,  $-\nabla \cdot p \nabla u = f$ ,  $p = p(x,y) \geq p_0 > 0$ . In particular, for the two-grid scheme we reobtain the basic result  $\rho \leq \frac{1}{2}(1+Kh)$  in the stronger form  $\|MG\|_{L_h} \leq \frac{1}{2}(1+Kh)$ . Computational results indicate that other constant coefficient results carry over as well.

*Additional keywords: algorithms; linear algebraic equations; problem solving; experimental data.*

Accession For	
NTIS	<input checked="" type="checkbox"/>
AD	<input type="checkbox"/>
AN	<input type="checkbox"/>
Availability Codes	
Dist	Special
A-1	



## I. Introduction

Multigrid methods are proving themselves as (very) successful tools for the solution of the algebraic equations associated with discretization of Elliptic Boundary-Value problems - see [1], [3], [4], [5], [9]. Nevertheless, it seems we are just beginning to "understand" this powerful idea. Hence, there is a need for continued probing, experimentation and new proofs - less for the sake of proof and more for the sake of insight.  $\rho^2 - 1$

In [2] Braess proposed and analyzed a class of multigrid methods. In particular, he considered a particular algorithm for the Poisson Equation - "Algorithm 2.1". He shows that the contraction number  $\rho$  for a two-grid method is given by

$$(1) \quad \rho \leq \frac{1}{2} !$$

This result holds whenever  $\Omega$  is a polygonal domain whose sides have slope  $\pm 1, 0$  or  $\infty$  and the discretization satisfies an additional condition (see  $\Omega I$  of section 2). In [8] Ries, Trottenberg and Winter discuss the class of MGR[v] methods for the Poisson Equation in a square. Using Fourier Analysis they obtain an explicit formula for the corresponding contraction numbers  $\rho[v]$ . In particular, they obtain - for two grids

$$(2) \quad \rho[0] = \frac{1}{2}, \quad \rho[1] = \frac{2}{27}, \quad \rho[v] = \frac{1}{2} \frac{(2v)^{2v}}{(2v+1)^{2v+1}} .$$

As it happens MGR[0] is the same as the "Algorithm 2.1" and the results of [2] and [8] are consistent. The results of [8] are more precise for more restricted problems.

In this report we consider the more general diffusion equation

$$(3) \quad \begin{aligned} -\nabla \cdot p(x,y) \nabla u &= f \quad \text{in } \Omega, \\ u &= 0 \quad \text{on } \partial\Omega \\ p(x,y) &\geq p_0 > 0 \quad \text{and} \end{aligned}$$

where  $\Omega$  may be a general bounded piecewise smooth domain, or,  $\Omega$  is a polygonal domain whose sides have slope  $\pm 1$ , 0, or  $\infty$ . We employ the usual five-point difference analog of (3) and seek to solve the (large) system of linear algebraic equations. We consider a class of linear multi-grid methods which include the MGR[v] methods when  $p(x,y) \equiv 1$ . Our basic result is the following: Consider the two-grid method. Then

$$\rho \leq \|MG\|_{L_h} \leq \frac{1}{2}(1+Kh)$$

where  $\| \cdot \|_{L_h}$  denotes the energy norm and  $K$  is a constant determined by  $p_0$  and  $\|\nabla p\|_{\infty}$ , the  $\infty$  norm of the gradient of  $p(x,y)$ . Moreover, the proof clearly indicates why one should expect great improvement when further "smoothing" is introduced.

Thus we extend the results of Braess [2], Riès, Trottenberg and Winter [8] to include a variable diffusion coefficient  $p(x,y)$  and more general regions.

In section 2 we formulate the problem and the basic two-grid method of solution. In section 3 we prove the basic estimate. This proof proceeds from a fundamental insight of McCormick and Ruge [7]. Section 4 describes the results of some computational experiments which lead one to believe that the results of [2.] are essentially correct for the variable coefficient case as well. These computations were carried out on the CRAY I at the Los Alamos National Laboratory. Finally, an appendix gives the basic "energy" estimate required in section 3.



## 2. The Problem

Given a (small) value  $h > 0$  let  $\{(x_k, y_j) = (kh, jh); k, j = 0, \pm 1, \pm 2, \dots\}$  be the associated mesh points in the  $x - y$  plane. Let

$$(2.1a) \quad R_E := \{(x_k, y_j); k+j \equiv 0 \pmod{2}\},$$

$$(2.1b) \quad R_O := \{(x_k, y_j); (k+j) \equiv 1 \pmod{2}\}.$$

Let  $\Omega$  be a bounded domain in the plane with a piecewise smooth boundary  $\partial\Omega$ . We wish to define the set of "interior" mesh points,  $\Omega_h$ . We assume that  $h$  is less than  $\frac{1}{4}$  the length of each smooth section of  $\partial\Omega$ .

The main result, (Theorem 3.1) for the two-grid iterative schemes is valid in quite general domains provided that we use a modification of "approximation of degree 0" (see [6]) to describe the boundary conditions

### Definition:\*

- (i) If  $(x_k, y_j) \in R_O \cap \Omega$  we say that  $(x_k, y_j) \in \Omega_h$  if the four neighbors  $\{(x_{k+1}, y_j), (x_{k-1}, y_j), (x_k, y_{j-1}), (x_k, y_{j+1})\}$  and the line segments from  $(x_k, y_j)$  to each of its neighbors all lie in  $\bar{\Omega}$ , the closure of  $\Omega$ .
- (ii) If  $(x_k, y_j) \in R_E \cap \Omega$  we say that  $(x_k, y_j) \in \Omega_h$  if the eight neighbors  $\{(x_{k+1}, y_j), (x_{k-1}, y_j), (x_k, y_{j-1}), (x_k, y_{j+1}), (x_{k+1}, y_{j+1}), (x_{k+1}, y_{j-1}), (x_{k-1}, y_{j+1}), (x_{k-1}, y_{j-1})\}$  and the line segments from  $(x_k, y_j)$  to each of its neighbors all lie in  $\bar{\Omega}$ .

---

(\*) We must consider the line segments from  $(x_k, y_j)$  to the neighbors only in the case of reentrant corners or cusps.

When  $\partial\Omega$  has a cusp or a corner at a point  $(x,y)$  we require that

$$(x,y) = (x_k, y_j) \in R_E.$$

The points  $(x_k, y_j) \in \overline{\Omega}/\Omega_h$  are the boundary points of  $\Omega_h$ . That is

$$\partial\Omega_h := \{(x_k, y_j) \in \overline{\Omega}/\Omega_h\}.$$

A true multigrid requires the use of many coarser grids. In such general regions the treatment of the boundary conditions on succeeding coarser grids gets complicated. In truth, the multigrid literature has barely touched on this question. In the case studied by Braess [2],  $\Omega$  is a polygonal domain whose sides have slope  $\pm 1, 0$  or  $\infty$  and the corners all belong to the coarsest (and hence, the finest) grid. For this case we note that (see Figure 1):

$$(\Omega I.a) \quad \partial\Omega_h \subset \partial\Omega$$

and

( $\Omega I.b$ ) if  $\sigma$  is a side of  $\Omega$  with slope  $\pm 1$ , then all the points of  $\partial\Omega_h$  which also lie on  $\partial\Omega$  belong to  $R_E$ .


For any function  $F(x,y)$  defined on the  $(x,y)$  plane we write

$$(2.2a) \quad F_{k,j} := F(x_k, y_j),$$

$$(2.2b) \quad F_{k+\frac{1}{2},j} := F((k+\frac{1}{2})h, y_j),$$

$$(2.2c) \quad F_{k,j+\frac{1}{2}} := F(x_k, (j+\frac{1}{2})h).$$

To obtain an approximate solution of (3) we seek a grid function  $\{u_{kj}\}$  defined on the mesh points and satisfying the system of equations:  
for  $(x_k, y_j) \in \Omega_h$



■ points of  $R_E \cap \overline{\Omega}$ ,  
Key  
 ○ points of  $R_0 \cap \overline{\Omega}$ .

Figure 1

$$(2.3a) \quad \begin{cases} \frac{1}{h^2} \{p_{k-\frac{1}{2},j}[U_{k,j}-U_{k-1,j}] - p_{k+\frac{1}{2},j}[U_{k+1,j}-U_{k,j}]\} + \\ \frac{1}{h^2} \{p_{k,j-\frac{1}{2}}[U_{k,j}-U_{k,j-1}] - p_{k,j+\frac{1}{2}}[U_{k,j+1}-U_{k,j}]\} = f_{kj}, \end{cases}$$

and, for  $(x_k, y_j) \notin \Omega_h$

$$(2.3b) \quad U_{kj} = 0.$$

We rewrite (2.3) as

$$(2.4a) \quad [L_h U]_{kj} = f_{kj}, \quad (x_k, y_j) \in \Omega_h,$$

$$(2.4b) \quad U_{kj} = 0, \quad (x_k, y_j) \notin \Omega_h.$$

We turn to the question of the solution of these linear algebraic equations via a "two-grid" method. Let

$$(2.5) \quad \Omega_E := R_E \cap \Omega_h, \quad \Omega_0 := R_0 \cap \Omega_h.$$

Our two grids are  $\Omega_h$  and  $\Omega_E$ . Let  $S_h$  and  $S_E$  be the spaces of grid functions defined on  $R_E \cup R_0$  and  $R_E$  which vanish outside  $\Omega_h$  and  $\Omega_E$ . Our first step is to set-up "communication" between these two spaces.

To be specific, we construct linear "interpolation" and "projection" operators  $I_h^E, I_E^h$  so that

$$(2.6a) \quad I_h^E: S_h \rightarrow S_E, \quad (\text{Projection}),$$

$$(2.6b) \quad I_E^h: S_E \rightarrow S_h. \quad (\text{Interpolation}).$$

Define the *interpolation operator*  $I_E^h$  by

$$(2.7a) \quad [I_E^h U]_{kj} = U_{kj}, \quad \text{if } (x_k, y_j) \in R_E,$$

and, if  $(x_k, y_j) \in \Omega_0$  then

$$(2.7b) \quad [I_E^h U]_{kj} = \frac{1}{c_{kj}} \{ p_{k-\frac{1}{2},j} U_{k-1,j} + p_{k+\frac{1}{2},j} U_{k+1,j} +$$

$$p_{k,j-\frac{1}{2}} U_{k,j-1} + p_{k,j+\frac{1}{2}} U_{k,j+1} \}$$

where

$$(2.7c) \quad c_{kj} = \{ p_{k+\frac{1}{2},j} + p_{k-\frac{1}{2},j} + p_{k,j-\frac{1}{2}} + p_{k,j+\frac{1}{2}} \}.$$

Finally, if  $(x_k, y_j) \in R_0/\Omega_0$  then (of course)

$$(2.7d) \quad [I_E^h U]_{kj} = 0.$$

Observe that (2.7a) implies that  $I_E^h$  is of full rank, i.e.,

$$\dim \text{Range } I_E^h = \dim S_E.$$

The *projection operator*  $I_h^E$  is defined by

$$(2.8) \quad I_h^E = \frac{1}{2} (I_E^h)^T.$$

Let

$$(2.9) \quad R := \text{Range } I_E^h.$$

The choice of interpolation operator  $I_E^h$  enables us to characterize  $R$  as follows:

Lemma 2.1: Let  $I_E^h$  be defined by (2.7). Then, a function  $U = U(h) \in S_h$  is in  $R$  if and only if

$$(2.10) \quad [L_h U]_{kj} = 0 \quad \forall (k,j) \text{ with } (x_k, y_j) \in \Omega_0.$$

We are now ready to describe the two-grid methods. Let  $G$  be a smoothing operator. That is, given  $u^0 \in S_h$  we construct  $\tilde{u}$  via

$$(2.11a) \quad \tilde{u} = Gu^0 = u^0 + B(f - L_h u^0) = G_0 u^0 + Bf$$

$$(2.11b) \quad G_0 = (I - BL_h)$$

where  $B$  is a given matrix and

$$(2.11c) \quad \|G_0\|_{L_h}^2 = \sup_{\psi \neq 0} \frac{\langle L_h G_0 \psi, G_0 \psi \rangle}{\langle L_h \psi, \psi \rangle} \leq 1.$$

Algorithm 2.1:

Step 1: Given  $u^0 \in S_h$ , form  $\tilde{u} = Gu^0$ .

Step 2: Construct  $\hat{u}$  via

$$\hat{u}_{kj} = \tilde{u}_{kj}, \quad (x_k, y_j) \in R_E$$

$$[L_h \hat{u}]_{kj} = f_{kj}, \quad (x_k, y_j) \in \Omega_0$$

$$\hat{u}_{kj} = 0, \quad (x_k, y_j) \in R_0/\Omega_0$$

That is: "relax" the equations on the "odd" points.

Step 3: Set  $r = f - L_h \hat{u}$ ,  $r_E = I_h^E r$ .

Step 4: Solve  $L_E \phi = r_E$  where  $L_E$  is the "coarse grid operator" to be described later.

Step 5: Set  $u^1 = \hat{u} + I_E^h \phi$ .

Step 6: Set  $u^1 \rightarrow u^0$  and return to step 1.

An important smoother  $G$  is the odd-even Gauss-Seidel scheme. That is, define  $H^0$  - relaxation on the odd points as in Step 2 -

$$(2.12a) \quad (H^0 u)_{kj} = u_{kj}, \quad (x_k, y_j) \in R_E,$$

and

$$(2.12b) \quad [L_h(H^0 u)]_{kj} = f_{kj}, \quad (x_k, y_j) \in \Omega_0,$$

$$(2.12c) \quad u_{k,j} = 0, \quad (x_k, y_j) \in R_0/\Omega_0.$$

Similarly define  $H^E$ , relaxation on the even points, by

$$(2.13a) \quad (H^E u)_{kj} = u_{kj}, \quad (x_k, y_j) \in R_0 \cup R_E/\Omega_E,$$

$$(2.13b) \quad [L_h(H^E u)]_{kj} = f_{kj}, \quad (x_k, y_j) \in \Omega_E.$$

Let  $v \geq 0$  be a integer. We obtain the generalized MGR[v] two-grid iterative scheme by choosing

$$(2.14) \quad G = (H^E H^0)^v.$$

We now describe two choices of the coarse grid operator  $L_E$ .

Case 1: Let

$$(2.15a) \quad a_{k-\frac{1}{2}, j-\frac{1}{2}} = \frac{1}{h^2} \left[ \frac{p_{k-\frac{1}{2}, j} p_{k-1, j-\frac{1}{2}}}{c_{k-1, j}} + \frac{p_{k, j-\frac{1}{2}} p_{k-\frac{1}{2}, j-1}}{c_{k, j-1}} \right],$$

$$(2.15b) \quad b_{k+\frac{1}{2}, j-\frac{1}{2}} = \frac{1}{h^2} \left[ \frac{p_{k, j-\frac{1}{2}} p_{k+\frac{1}{2}, j-1}}{c_{k, j-1}} + \frac{p_{k+\frac{1}{2}, j} p_{k+1, j-\frac{1}{2}}}{c_{k+1, j}} \right],$$

$$(2.15c) \quad d_{kj} = [a_{k-\frac{1}{2}, j-\frac{1}{2}} + a_{k+\frac{1}{2}, j+\frac{1}{2}} + b_{k+\frac{1}{2}, j-\frac{1}{2}} + b_{k-\frac{1}{2}, j+\frac{1}{2}}].$$

Then, if  $(k+j) \equiv 0 \pmod{2}$ .

$$(2.16) \quad [L_E^{(1)} U]_{kj} = -a_{k+\frac{1}{2}, j+\frac{1}{2}} U_{k+1, j+1} - a_{k-\frac{1}{2}, j-\frac{1}{2}} U_{k-1, j-1} \\ - b_{k+\frac{1}{2}, j-\frac{1}{2}} U_{k+1, j-1} - b_{k-\frac{1}{2}, j+\frac{1}{2}} U_{k-1, j+1} + d_{kj} U_{kj}.$$

Case 2: (The Standard Case): if  $k+j \equiv 0 \pmod{2}$  then

$$(2.17a) \quad [L_E^{(2)} U]_{kj} = \frac{1}{2h^2} \{-p_{k+\frac{1}{2}, j+\frac{1}{2}} U_{k+1, j+1} - p_{k+\frac{1}{2}, j-\frac{1}{2}} U_{k+1, j-1} \\ - p_{k-\frac{1}{2}, j-\frac{1}{2}} U_{k-1, j-1} - p_{k-\frac{1}{2}, j+\frac{1}{2}} U_{k-1, j+1} + S_{kj} U_{kj}\}$$

where

$$(2.17b) \quad S_{kj} = \{p_{k+\frac{1}{2}, j+\frac{1}{2}} + p_{k+\frac{1}{2}, j-\frac{1}{2}} + p_{k-\frac{1}{2}, j+\frac{1}{2}} + p_{k-\frac{1}{2}, j-\frac{1}{2}}\}.$$



### 3. Analysis of the Algorithm

We begin our analysis with an observation which is (by now) well known among multigrid theorists (see [7]). Let

$$(3.1) \quad \hat{L}_E := I_h^E L_h I_E^h .$$

Consider Steps 4-5 of the two-grid iteration. Suppose we replace  $L_E$  by  $\hat{L}_E$ , i.e., suppose we find the function  $\psi$  which satisfies

$$\hat{L}_E \psi = r_E ,$$

and set

$$u^1 = \hat{u} + I_E^h \psi .$$

We claim that

$$L_h u^1 = f ,$$

i.e.  $u^1$  is the desired solution! To see this we set

$$(3.2a) \quad \hat{\epsilon} = U - \hat{u}$$

and observe that Step 2 implies that if  $k + j \equiv 1 \pmod{2}$ , then

$$(L_h \hat{\epsilon})_{kj} = (L_h U - L_h \hat{u})_{kj} = (f - L_h \hat{u})_{kj} = 0 .$$

Hence Lemma 2.1 asserts that there is a function  $V \in S_E$  and

$$(3.2b) \quad \hat{\epsilon} = I_E^h V .$$

We now verify that

$$\hat{L}_E V = I_h^E (L_h I_E^h V) = I_h^E L_h \hat{\varepsilon} = r_E.$$

Hence,

$$\psi = V$$

and

$$(3.3) \quad \hat{u} - I_E^h \psi = \hat{u} - \hat{\varepsilon} = U!!$$

Unfortunately we have chosen Step 4 with  $L_E$  and not  $\hat{L}_E$ . This choice was not merely pique on our part (or the part of Braes and Ries, Trottenberg and Winter). The point is -- having chosen  $L_E$  as a five point star we can now proceed to replace Step 4 with a new two grid step -- i.e. we can build a true multigrid.

In any case, the problem of Step 4 is seen to be

$$L_E \phi = \hat{L}_E \psi,$$

where, as we see from Lemma 2.1,  $I_E^h \psi$  is the  $L_h$  projection of  $\tilde{\varepsilon}$  into  $R$ . Hence

$$(3.4a) \quad \|\hat{\varepsilon}\|_{L_h} = \|I_E^h \psi\|_{L_h} \leq \|\tilde{\varepsilon}\|_{L_h}.$$

We will give a complete description of  $\hat{L}_E$  in the appendix. For now, we write

$$(3.5) \quad \hat{L}_E = \frac{1}{2} L_E + \frac{1}{2} \tilde{L}_E$$

where  $\tilde{L}_E$  is defined by this equation. Observe that both  $\hat{L}_E$  and  $L_E$  (either  $L_E^{(1)}$  or  $L_E^{(2)}$ ) are symmetric, positive definite operators.

Hence the associated  $\tilde{L}_E$  is a symmetric operator. Our main estimate is

Lemma 3.1: For  $L_E = L_E^{(1)}$  or  $L_E = L_E^{(2)}$ , there is a constant  $K$ , depending only on  $\|\nabla p\|_\infty$ , the maximum norm of the first derivatives of the diffusion coefficient  $p(x,y)$ , and  $p_0$  such that, for all  $\phi \in S_E$ ,  $\phi \neq 0$  we have

$$(3.6) \quad -Kh \leq \frac{\langle \tilde{L}_E \phi, \phi \rangle}{\langle L_E \phi, \phi \rangle} \leq 2(1+Kh) .$$

Proof: See Theorem A of the Appendix.

Consider the eigenvalue problem

$$(3.7a) \quad (\lambda L_E - \hat{L}_E) \psi = 0, \quad \psi \neq 0$$

which is equivalent to

$$(3.7b) \quad (\lambda I - L_E^{-1} \hat{L}_E) \psi = 0, \quad \psi \neq 0 .$$

Using (3.5) we see that this problem is equivalent to

$$(3.8a) \quad [(2\lambda - 1)L_E - \tilde{L}_E] \psi = 0 .$$

From Lemma 3.1 we find

$$(3.9) \quad \frac{1-Kh}{2} \leq \lambda \leq \frac{3+2Kh}{2} .$$

Theorem 3.1: Let

$$\epsilon^0 = U - u^0, \quad \epsilon^1 = U - u^1$$

then

$$(3.10) \quad \|\epsilon^1\|_{L_h} \leq \frac{1}{2}(1+Kh) \|\epsilon^0\|_{L_h} .$$

Proof: We have

$$\tilde{\epsilon} = U - u^0 = G_0 \epsilon^0 .$$

Using (2.11c) we see that

$$(3.11) \quad \|\tilde{\epsilon}\|_{L_h} \leq \|\epsilon^0\|_{L_h} .$$

From (3.2a), (3.3b) and Step 5 of the multigrid algorithm we have

$$(3.12a) \quad \epsilon^1 = \hat{\epsilon} - I_E^h \phi = I_E^h (\psi - \phi) ,$$

and

$$(3.12b) \quad L_E \phi = \hat{L}_E \psi .$$

Hence

$$(3.13) \quad \psi - \phi = (I - L_E^{-1} \hat{L}_E) \psi .$$

Thus

$$\begin{aligned} (3.14a) \quad \langle L_h \epsilon^1, \epsilon^1 \rangle_h &= \langle L_h I_E^h (\psi - \phi), I_E^h (\psi - \phi) \rangle_h = 2 \langle I_h^E L_h I_E^h (\psi - \phi), (\psi - \phi) \rangle_E \\ &= 2 \langle \hat{L}_E (\psi - \phi), (\psi - \phi) \rangle_E \\ &= 2 \langle \hat{L}_E (I - L_E^{-1} \hat{L}_E) \psi, (I - L_E^{-1} \hat{L}_E) \psi \rangle_E , \end{aligned}$$

$$(3.14b) \quad \|\epsilon^1\|_{L_h}^2 = 2 \langle [I - \hat{L}_E^{-1/2} L_E^{-1} \hat{L}_E^{1/2}] \hat{L}_E^{1/2} \psi, [I - \hat{L}_E^{-1/2} L_E^{-1} \hat{L}_E^{1/2}] \hat{L}_E^{1/2} \psi \rangle_E .$$

Since the eigenvalues of  $L_E^{-1} \hat{L}_E$  are also the eigenvalues of the symmetric operator  $\hat{L}_E^{-1/2} L_E^{-1} \hat{L}_E^{1/2}$ , (3.9) implies that the eigenvalues  $\mu$  of the symmetric operator  $(I - \hat{L}_E^{-1/2} L_E^{-1} \hat{L}_E^{1/2})$  satisfy

$$-\frac{1}{2}(1+Kh) \leq \mu \leq \frac{1}{2}(1+Kh) .$$

Thus, (3.14b) implies

$$\begin{aligned}
 \| \epsilon^1 \|_{L_h}^2 &\leq 2\mu^2 \langle \hat{L}_E^{\frac{1}{2}} \psi, \hat{L}_E^{\frac{1}{2}} \psi \rangle_E = 2\mu^2 \langle \hat{L}_E \psi, \psi \rangle_E = 2\mu^2 \langle L_h I_E^h \psi, (I_h^E)^T \psi \rangle_h \\
 &= 2\mu^2 \frac{1}{2} \langle L_h I_E^h \psi, I_E^h \psi \rangle_h \\
 &= \mu^2 \langle L_h \hat{\epsilon}, \hat{\epsilon} \rangle_h = \mu^2 \| \hat{\epsilon} \|_{L_h}^2.
 \end{aligned}$$

This result, together with (3.4a) and (3.11) implies the Theorem.

#### 4. Experimental Results

In order to demonstrate that the results of section 3 are valid for the variable coefficient case an experimental project was undertaken. The essence of this project was to write a computer program which implemented Algorithm 2.1. By experimenting with different functions  $p(x,y)$  and different true solutions  $u(x,y)$  it was shown that formula (2) of section 1 is valid for the variable coefficient case. The region  $\Omega$  is the unit square.

The computer program runs in an interactive fashion and allows the user to provide a number of parameters. These include  $N$ , the number of points on a side of  $\Omega_h$ , the fine grid and  $v$ , the number of smoothing iterations. Starting with a particular initial guess, Algorithm 2.1 was then repeated until the discrete  $L_2$  norm of the residual was less than  $10^{-8}$ . For the initial guess  $U^0$ , interior points of  $\Omega_E$  were set to 5 while interior points of  $\Omega_0$  were set to -5.

Experiments were done with  $L_E$ , the coarse grid operator chosen to be both  $L_E^{(1)}$  and  $L_E^{(2)}$ . The calculation of  $L_E^{(1)}$  was complicated by the fact that for points of  $\Omega_E$  for which  $L_E^{(1)}$  refers to points of  $\partial\Omega_h$ , formula (2.15c) does not apply. The reason for this is because the computation of either  $a_{k\pm\frac{1}{2}j\pm\frac{1}{2}}$  or  $b_{k\pm\frac{1}{2}j\mp\frac{1}{2}}$  involves referring to points outside of  $\Omega_h$ . Of course since  $U_{kj} = 0$  if  $U_{kj} \in \partial\Omega_h$  we set  $a_{k\pm\frac{1}{2}j\pm\frac{1}{2}}$  and  $b_{k\pm\frac{1}{2}j\mp\frac{1}{2}}$  to zero when  $U_{k\pm\frac{1}{2}j\pm\frac{1}{2}}$  and  $U_{k\pm\frac{1}{2}j\mp\frac{1}{2}}$  are in  $\partial\Omega_h$ . However, we still need a value for  $d_{kj}$  for the two nearest interior points. For the four corners points, we set  $d_{kj}$  to be the value of  $d_{kj}$  of the nearest interior point. As the mesh gets finer, this approximation to the true

$d_{kj}$  improves. However, in almost all of the experiments the rate of convergence using  $L_E^{(1)}$  was not quite as good as the rate obtained using  $L_E^{(2)}$ .

The tables below list the functions  $p(x,y)$  and the true solutions  $u(x,y)$  used for the experiments. For each problem the numerical results obtained using both  $L_E^{(1)}$  and  $L_E^{(2)}$  are displayed.  $N$  corresponds to the number of interior points on a side of  $\Omega_h$  and  $\nu$  corresponds to the number of smoothing iterations. The smoother used was the odd-even Gauss-Seidel scheme as described in section 2.  $\sigma(\nu)$  in the tables corresponds to the theoretical rate given in equation (2) of section 1. The theoretical rate has only been proven to be valid, when  $\nu > 0$ , in the constant coefficient case. However, as can be seen from the numerical results it appears to be valid in the variable coefficient case as well.

In conclusion, the numerical results demonstrate the validity of Theorem 3.1 for the case  $\nu = 0$  and support extending equation (2) of section 1 to the variable coefficient case.

Table I, Experimental Results

Problem 1,  $p(x,y) = 1$ ,  $u(x,y) = 0$

$$L_E = L_E^{(1)}$$

$N \backslash v$	0	1	2	3
15	.4858	.0646	.0344	.0200
31	.4844	.0696	.0375	.0252
63	.4836	.0708	.0386	.0263
$\sigma(v)$	.5000	.0741	.0410	.0283

$$L_E = L_E^{(2)}$$

$N \backslash v$	0	1	2	3
15	.4858	.0646	.0344	.0200
31	.4844	.0696	.0375	.0252
63	.4836	.0708	.0386	.0263
$\sigma(v)$	.5000	.0741	.0410	.0283

Problem 2,  $p(x,y) = 1$ ,  $u(x,y) = \sin \pi x \sin \pi y$

$$L_E = L_E^{(1)}$$

$N \backslash v$	0	1	2	3
15	.4858	.0646	.0344	.0200
31	.4844	.0696	.0375	.0252
63	.4836	.0708	.0386	.0263
$\sigma(v)$	.5000	.0741	.0410	.0283

$$L_E = L_E^{(2)}$$

$N \backslash v$	0	1	2	3
15	.4858	.0696	.0344	.0200
31	.4844	.0696	.0375	.0252
63	.4836	.0708	.0386	.0263
$\sigma(v)$	.5000	.0741	.0410	.0283



Problem 3,  $p(x,y) = 1$ ,  $u(x,y) = x(1-x)y(1-y)$

$$L_E = L_E^{(1)}$$

N \ v	0	1	2	3
15	.4858	.0646	.0344	.0200
31	.4844	.0696	.0375	.0252
63	.4836	.0708	.0386	.0263
$\sigma(v)$	.5000	.0741	.0410	.0283

$$L_E = L_E^{(2)}$$

N \ v	0	1	2	3
15	.4858	.0646	.0344	.0200
31	.4844	.0696	.0375	.0252
63	.4836	.0708	.0386	.0263
$\sigma(v)$	.5000	.0741	.0410	.0283

Problem 4,  $p(x,y) = e^{xy}$ ,  $u(x,y) = xe^{xy} \sin \pi x \sin \pi y$

$$L_E = L_E^{(1)}$$

N \ v	0	1	2	3
15	.4863	.0760	.0437	.0292
31	.4841	.0742	.0425	.0303
63	.4842	.0720	.0401	.0283
$\sigma(v)$	.5000	.0741	.0410	.0283

$$L_E = L_E^{(2)}$$

N \ v	0	1	2	3
15	.4858	.0643	.0342	.0199
31	.4840	.0697	.0373	.0252
63	.4841	.0709	.0384	.0264
$\sigma(v)$	.5000	.0741	.0410	.0283

Problem 5,  $p(x,y) = \frac{1}{(3-x)(3-y)}$ ,  $u(x,y) = e^{xy} \sin \pi x \sin \pi y$

$$L_E = L_E^{(1)}$$

N \ v	0	1	2	3
15	.4841	.0708	.0393	.0270
31	.4819	.0713	.0398	.0276
63	.4820	.0709	.0386	.0268
$\sigma(v)$	.5000	.0741	.0410	.0283

$$L_E = L_E^{(2)}$$

N \ v	0	1	2	3
15	.4839	.0643	.0339	.0199
31	.4819	.0694	.0373	.0250
63	.4820	.0706	.0381	.0261
$\sigma(v)$	.5000	.0741	.0410	.0283

Problem 6,  $p(x,y) = e^x(1 + \frac{1}{2} \sin \pi y)$ ,  $u(x,y) = e^{xy} \sin \pi x \sin \pi y$

$$L_E = L_E^{(1)}$$

N \ v	0	1	2	3
15	.4879	.1084	.0727	.0565
31	.4854	.0901	.0582	.0442
63	.4851	.0784	.0473	.0350
$\sigma(v)$	.5000	.0741	.0410	.0283

$$L_E = L_E^{(2)}$$

N \ v	0	1	2	3
15	.4869	.0686	.0377	.0255
31	.4851	.0710	.0390	.0270
63	.4850	.0715	.0390	.0270
$\sigma(v)$	.5000	.0741	.0410	.0283

Problem 7,  $p(x,y) = e^{-xy}$ ,  $u(x,y) = (1-e^x)(x-1)y \cos \frac{\pi y}{2}$

$$L_E = L_E^{(1)}$$

N \ v	0	1	2	3
15	.4857	.0797	.0482	.0351
31	.4842	.0739	.0431	.0312
63	.4836	.0714	.0399	.0278
$\sigma(v)$	.5000	.0741	.0410	.0283

$$L_E = L_E^{(2)}$$

N \ v	0	1	2	3
15	.4853	.0650	.0347	.0207
31	.4841	.0697	.0376	.0253
63	.4835	.0708	.0386	.0263
$\sigma(v)$	.5000	.0741	.0410	.0283

Problem 8,  $p(x,y) = e^{(\sin \frac{\pi x}{2} \cos \pi y)}$ ,  $u(x,y) = e^{-xy} x(x-1)y(y-1)$

$$L_E = L_E^{(1)}$$

N \ v	0	1	2	3
15	.4849	.0772	.0451	.0296
31	.4839	.0751	.0437	.0313
63	.4842	.0721	.0404	.0289
$\sigma(v)$	.5000	.0741	.0410	.0283

$$L_E = L_E^{(1)}$$

N \ v	0	1	2	3
15	.4843	.0645	.0342	.0202
31	.4837	.0697	.0373	.0253
63	.4842	.0710	.0385	.0264
$\sigma(v)$	.5000	.0741	.0410	.0283

Problem 9,  $p(x,y) = \begin{bmatrix} 1 & \text{for } \frac{1}{4} \leq x \leq \frac{3}{4}, \quad \frac{3}{4}-x \leq y \leq \frac{1}{4}+x, \quad x-\frac{1}{4} \leq y \leq \frac{5}{4}-x \\ 6 & \text{otherwise} \end{bmatrix}$

$$u(x,y) = e^{-xy} x(x-1)y(y-1)$$

$$L_E = L_E^{(1)}$$

$N \backslash v$	0	1	2	3
15	.4848	.0648	.0311	.0174
31	.4857	.0698	.0377	.0248
63	.4855	.0711	.0389	.0266
$\sigma(v)$	.5000	.0741	.0410	.0283

$$L_E = L_E^{(2)}$$

$N \backslash v$	0	1	2	3
15	.4649	.1815	.1015	.0634
31	.4854	.1186	.1291	.1046
63	.4855	.0712	.0393	.0275
$\sigma(v)$	.5000	.0741	.0410	.0283

### Appendix

In this section we determine  $\hat{L}_E$  and the quadratic forms

$$\langle \hat{L}_E \psi, \psi \rangle, \quad \langle \tilde{L}_E \psi, \psi \rangle, \quad \langle L_E \psi, \psi \rangle$$

Let  $U \in S_E$ , let  $(x_k, y_j) \in \Omega_E$ . Then

$$(A.1) \quad [\hat{L}_E U]_{kj} = \frac{1}{2} [L_h I_E^h U]_{kj}.$$

For any  $V \in S_h$ ,  $[L_h V]_{kj}$  involves the four values  $V_{k\pm 1, j}$ ,  $V_{k, j\pm 1}$ .

Therefore we consider the four squares I, II, III, IV (see fig. 2) with vertices

$$(A.2a) \quad \text{I: } \{(x_k, y_j), (x_{k+1}, y_{j+1}), (x_{k+2}, y_j), (x_{k+1}, y_{j-1})\},$$

$$(A.2b) \quad \text{II: } \{(x_k, y_k), (x_{k+1}, y_{j+1}), (x_k, y_{j+2}), (x_{k-1}, y_{j+1})\},$$

$$(A.2c) \quad \text{III: } \{(x_k, y_j), (x_{k-1}, y_{j+1}), (x_{k-2}, y_j), (x_{k-1}, y_{j-1})\},$$

$$(A.2d) \quad \text{IV: } \{(x_k, y_j), (x_{k-1}, y_{j-1}), (x_k, y_{j-2}), (x_{k+1}, y_{j-1})\}.$$

In each square the value of  $[I_E^h U]$  at the center point is a weighted average (given by (2.7b), (2.7c)) of the values of  $U$  at the corners. Thus, in general  $\hat{L}_E$  is a 9-point operator based on the 9 vertices of these four squares.

Since  $\hat{L}_E$  is a symmetric operator it takes the form

$$(A.3) \quad \begin{aligned} [\hat{L}_E U]_{kj} = & E_{kj} U_{kj} - \alpha_{k+1, j} U_{k+2, j} - \alpha_{k-1, j} U_{k-2, j} \\ & - \beta_{k, j+1} U_{k, j+2} - \beta_{k, j-1} U_{k, j-2} \\ & - \gamma_{k+\frac{1}{2}, j+\frac{1}{2}} U_{k+1, j+1} - \gamma_{k-\frac{1}{2}, j-\frac{1}{2}} U_{k-1, j-1} \\ & - \sigma_{k-\frac{1}{2}, j+\frac{1}{2}} U_{k-1, j+1} - \sigma_{k+\frac{1}{2}, j-\frac{1}{2}} U_{k+1, j-1}. \end{aligned}$$




Figure 2

Lemma A.1: Let

$$(A.4a) \quad E_{kj} = E_{kj}^0 + \tilde{E}_{kj}$$

where

$$(A.4b) \quad \begin{aligned} E_{kj}^0 = & \alpha_{k+1,j} + \alpha_{k-1,j} + \beta_{k,j-1} + \beta_{k,j+1} + \gamma_{k+\frac{1}{2},j+\frac{1}{2}} \\ & + \gamma_{k-\frac{1}{2},j-\frac{1}{2}} + \sigma_{k+\frac{1}{2},j-\frac{1}{2}} + \sigma_{k-\frac{1}{2},j+\frac{1}{2}} . \end{aligned}$$

Then

$$(A.5) \quad \begin{aligned} \langle \hat{L}_E^{\psi,\psi} \rangle = & \sum \alpha_{k+1,j} [\psi_{k+2,j} - \psi_{kj}]^2 \\ & + \sum \beta_{k,j+1} [\psi_{k,j+2} - \psi_{kj}]^2 + \sum \gamma_{k+\frac{1}{2},j+\frac{1}{2}} [\psi_{k+1,j+1} - \psi_{kj}]^2 \\ & + \sum \sigma_{k+\frac{1}{2},j-\frac{1}{2}} [\psi_{k+1,j-1} - \psi_{kj}]^2 + \sum \tilde{E}_{kj} \psi_{kj}^2 . \end{aligned}$$

Proof: Summation by parts.

Similar calculations yield

Lemma A.2: Using the definitions (2.15), (2.17) we have

$$(A.6) \quad \begin{aligned} \langle L_E^{(1)} \psi, \psi \rangle = & \sum a_{k+\frac{1}{2},j+\frac{1}{2}} [\psi_{k+1,j+1} - \psi_{kj}]^2 \\ & + \sum b_{k+\frac{1}{2},j-\frac{1}{2}} [\psi_{k+1,j-1} - \psi_{kj}]^2 , \end{aligned}$$

and

$$(A.7) \quad \begin{aligned} \langle L_E^{(2)} \psi, \psi \rangle = & \frac{1}{2h^2} \sum p_{k+\frac{1}{2},j+\frac{1}{2}} [\psi_{k+1,j+1} - \psi_{kj}]^2 \\ & + \frac{1}{2h^2} \sum p_{k+\frac{1}{2},j-\frac{1}{2}} [\psi_{k+1,j-1} - \psi_{kj}]^2 . \end{aligned}$$

We now compute the contribution of each square to  $[\hat{L}_E^U]_{kj}$  and the two quadratic forms. In evaluating  $[L_h I_E^h U]_{kj}$  we have five terms. The four terms

$$-p_{k\pm\frac{1}{2},j}(I_E^h U)_{k\pm 1,j}, \quad -p_{k,j\pm\frac{1}{2}}(I_E^h U)_{k,j\pm 1}$$

are clearly associated with squares  $\begin{pmatrix} I \\ III \end{pmatrix}$  and  $\begin{pmatrix} II \\ IV \end{pmatrix}$  respectively. It is convenient to agree that

$p_{k\pm\frac{1}{2},j} U_{kj}$  is associated with square  $\begin{pmatrix} I \\ III \end{pmatrix}$ ,  
and  
 $p_{k,j\pm\frac{1}{2}} U_{kj}$  is associated with square  $\begin{pmatrix} II \\ IV \end{pmatrix}$ .

Let  $E_{kj}(R)$ ,  $\alpha_{k\pm 1,j}(R)$ ,  $\beta_{k,j\pm 1}(R)$ ,  $\gamma_{k\pm\frac{1}{2},j\pm\frac{1}{2}}(R)$ ,  $\sigma_{k\pm\frac{1}{2},j\mp\frac{1}{2}}(R)$  denote the contributions of square  $R$  to the corresponding coefficients  $E_{kj}$ ,  $\alpha_{k\pm 1,j}$ ,  $\beta_{k,j\pm 1}$ ,  $\gamma_{k\pm\frac{1}{2},j\pm\frac{1}{2}}$ ,  $\sigma_{k\pm\frac{1}{2},j\mp\frac{1}{2}}$  of  $\hat{L}_E$ .

Consider square I. We must consider two cases, either  $(x_{k+1}, y_j) \in \Omega_h$  or  $(x_{k+1}, y_j) \notin \Omega_h$ . The following geometric lemma is essential to understanding the computations in the latter case.

Lemma A.3: Suppose  $(x_{k+1}, y_j) \notin \Omega_h$ . Then either  $(x_{k+2}, y_j) \notin \bar{\Omega}$  or the line segment from  $(x_{k+1}, y_j)$  to  $(x_{k+2}, y_j)$  is not entirely in  $\bar{\Omega}$ . Further  $(x_{k+1}, y_{j+1}), (x_{k+1}, y_{j-1}) \notin \Omega_E$ .

Proof: (see Figure 3). Since  $(x_k, y_j) \in \Omega_E$  the points  $(x_{k+1}, y_{j+1}), (x_{k+1}, y_j), (x_{k+1}, y_{j-1}) \in \bar{\Omega}$ . On the other hand,  $(x_{k+1}, y_j) \notin \Omega_h$  implies that either  $(x_{k+2}, y_j) \notin \bar{\Omega}$  or the line segment from  $(x_{k+1}, y_j)$  to  $(x_{k+2}, y_j) \notin \bar{\Omega}$ . If  $(x_{k+2}, y_j) \notin \bar{\Omega}$  then clearly  $(x_{k+1}, y_{j+1})$  and  $(x_{k+1}, y_{j-1}) \notin \Omega_E$ .




Figure 3



If  $(x_{k+2}, y_j) \in \bar{\Omega}$  then a portion of  $\partial\Omega$  crosses the line segment  $x_{k+1} \leq x < x_{k+2}$ ,  $y = y_j$ . If that portion of the boundary continues smoothly near  $(x_{k+1}, y_j)$ , then the line segments from  $(x_{k+1}, y_{j\pm 1})$  to  $(x_{k+2}, y_j)$  are not entirely in  $\bar{\Omega}$ . Finally, if there is a non-convex corner  $(x, y)$  near  $(x_{k+1}, y_j)$  that corner  $(x, y) \in R_E$ . Hence that corner must be  $(x_{k+1}, y_{j+1})$  or  $(x_{k+1}, y_{j-1})$  which is therefore not in  $\Omega_E$ . The other one is not in  $\Omega_E$  because  $h$  is less than  $\frac{1}{4}$  the length of smooth segments of  $\partial\Omega$ .

We return to the calculation of  $E_{kj}(I)$ ,  $\alpha_{k+1,j}(I)$ ,  $\gamma_{k+\frac{1}{2},j+\frac{1}{2}}(I)$ , and  $\sigma_{k+\frac{1}{2},j-\frac{1}{2}}(I)$ . Square I does not contribute to the other coefficients.

Case 1:  $(x_{k+1}, y_j) \in \Omega_h$ :

A straight forward calculation yields

$$(A.8a) \quad E_{kj}(I) = \frac{1}{2h^2} \left[ p_{k+\frac{1}{2},j} - \frac{(p_{k+\frac{1}{2},j})^2}{c_{k+1,j}} \right],$$

$$(A.8b) \quad \alpha_{k+1,j}(I) = \frac{1}{2h^2} \frac{p_{k+\frac{1}{2},j} p_{k+\frac{3}{2},j}}{c_{k+1,j}},$$

$$(A.8c) \quad \gamma_{k+\frac{1}{2},j+\frac{1}{2}}(I) = \frac{1}{2h^2} \frac{p_{k+\frac{1}{2},j} p_{k+1,j+\frac{1}{2}}}{c_{k+1,j}},$$

$$(A.8d) \quad \sigma_{k+\frac{1}{2},j-\frac{1}{2}}(I) = \frac{1}{2h^2} \frac{p_{k+\frac{1}{2},j} p_{k+1,j-\frac{1}{2}}}{c_{k+1,j}}.$$

Case 2:  $(x_{k+1}, y_j) \notin \Omega_h$ :

In this case we set

$$(A.9a) \quad E_{kj}(I) = \frac{1}{2h^2} p_{k+\frac{1}{2},j},$$

$$(A.9b) \quad \gamma_{k+\frac{1}{2},j+\frac{1}{2}}(I) = \frac{1}{2h^2} \frac{p_{k+\frac{1}{2},j} p_{k+1,j+\frac{1}{2}}}{c_{k+1,j}},$$

$$(A.9c) \quad \sigma_{k+\frac{1}{2},j-\frac{1}{2}}(I) = \frac{1}{2h^2} \frac{p_{k+\frac{1}{2},j} p_{k+1,j-\frac{1}{2}}}{c_{k+1,j}}.$$

$$(A.9d) \quad \alpha_{k+1,j}(I) = E_{kj}(I) - \gamma_{k+\frac{1}{2},j+\frac{1}{2}}(I) - \sigma_{k+\frac{1}{2},j-\frac{1}{2}}(I).$$

Observe that  $\alpha_{k+1,j} > 0$ , and since  $u_{k+2,j} = u_{k+1,j+1} = u_{k+1,j-1} = 0$ , the choices of  $\alpha_{k+1,j}(I)$ ,  $\gamma_{k+\frac{1}{2},j+\frac{1}{2}}(I)$  and  $\sigma_{k+\frac{1}{2},j-\frac{1}{2}}(I)$  do not effect the value of  $\hat{L}_E$ .

Consider Square II.

Case 1:  $(x_k, y_{j+1}) \in \Omega_h$ :

In this case we obtain

$$(A.10a) \quad E_{kj}(II) = \frac{1}{2h^2} \left[ p_{k,j+\frac{1}{2}} - \frac{(p_{k,j+\frac{1}{2}})^2}{c_{k,j+1}} \right],$$

$$(A.10b) \quad \beta_{k,j+1}(II) = \frac{1}{2h^2} \frac{p_{k,j+\frac{1}{2}} p_{k,j+\frac{3}{2}}}{c_{k,j+1}},$$

$$(A.10c) \quad \gamma_{k+\frac{1}{2}, j+\frac{1}{2}}(II) = \frac{1}{2h^2} \frac{p_{k, j+\frac{1}{2}} p_{k+\frac{1}{2}, j+1}}{c_{k, j+1}},$$

$$(A.10d) \quad \sigma_{k-\frac{1}{2}, j+\frac{1}{2}}(II) = \frac{1}{2h^2} \frac{p_{k, j+\frac{1}{2}} p_{k-\frac{1}{2}, j+1}}{c_{k, j+1}},$$

Case 2:  $(x_k, y_{j+1}) \notin \Omega_h$ :

Using arguments similar to those used in case 2 of square I we have

$$(A.11a) \quad E_{kj}(II) = \frac{1}{2h^2} p_{k, j+\frac{1}{2}}.$$

As for  $\gamma_{k+\frac{1}{2}, j+\frac{1}{2}}(II)$  and  $\sigma_{k-\frac{1}{2}, j+\frac{1}{2}}(II)$  we may use the formulae of (A.10c) and (A.10d). Finally

$$(A.11b) \quad \beta_{k, j+1}(II) = E_{kj}(II) - \gamma_{k+\frac{1}{2}, j+\frac{1}{2}}(II) - \sigma_{k-\frac{1}{2}, j+\frac{1}{2}}(II).$$

Because  $\hat{L}_E$  is symmetric it is not necessary to compute the contributions from squares III and IV. We now make a similar decomposition of the coefficients of  $L_E^{(1)}$ . Set

$$(A.12a) \quad a_{k+\frac{1}{2}, j+\frac{1}{2}}(I) = \frac{1}{h^2} \frac{p_{k+\frac{1}{2}, j} p_{k+1, j+\frac{1}{2}}}{c_{k+1, j}},$$

$$(A.12b) \quad a_{k+\frac{1}{2}, j+\frac{1}{2}}(II) = \frac{1}{h^2} \frac{p_{k, j+\frac{1}{2}} p_{k+\frac{1}{2}, j+1}}{c_{k, j+1}},$$

$$(A.12c) \quad b_{k+\frac{1}{2}, j-\frac{1}{2}}(I) = \frac{1}{h^2} \frac{p_{k+\frac{1}{2}, j} p_{k+1, j-\frac{1}{2}}}{c_{k+1, j}},$$

$$(A.12d) \quad b_{k-\frac{1}{2}, j+\frac{1}{2}}(II) = \frac{1}{h^2} \frac{p_{k, j+\frac{1}{2}} p_{k-\frac{1}{2}, j+1}}{c_{k, j+1}}.$$

$$(A.13) \quad d_{kj}(I) = a_{k+\frac{1}{2}, j+\frac{1}{2}}(I) + b_{k+\frac{1}{2}, j-\frac{1}{2}}(I).$$

Let  $\langle \hat{L}_E U, U \rangle_I$  and  $\langle L_E^{(1)} U, U \rangle_I$  denote the contribution of square I to the quadratic forms  $\langle \hat{L}_E U, U \rangle$  and  $\langle L_E^{(1)} U, U \rangle$  respectively. Then, using lemma A.1 we see that

$$(A.11a) \quad \left\{ \begin{aligned} \langle \hat{L}_E \psi, \psi \rangle_I &= \alpha_{k+1, j}(I) [\psi_{k+2, j} - \psi_{k, j}]^2 + \\ &\beta_{k+1, j}(I) [\psi_{k+1, j+1} - \psi_{k+1, j-1}]^2 + \gamma_{k+\frac{1}{2}, j+\frac{1}{2}}(I) [\psi_{k+1, j+1} - \psi_{k, j}]^2 \\ &+ \gamma_{k+\frac{3}{2}, j-\frac{1}{2}}(I) [\psi_{k+2, j} - \psi_{k+1, j-1}]^2 + \sigma_{k+\frac{1}{2}, j-\frac{1}{2}}(I) [\psi_{k+1, j-1} - \psi_{k, j}]^2 \\ &+ \sigma_{k+\frac{3}{2}, j+\frac{1}{2}}(I) [\psi_{k+1, j+1} - \psi_{k+2, j}]^2, \end{aligned} \right.$$

and

$$(A.11b) \quad \left\{ \begin{aligned} \langle L_E^{(1)} \psi, \psi \rangle_I &= a_{k+\frac{1}{2}, j+\frac{1}{2}}(I) [\psi_{k+1, j+1} - \psi_{k, j}]^2 \\ &+ a_{k+\frac{3}{2}, j-\frac{1}{2}}(I) [\psi_{k+2, j} - \psi_{k+1, j-1}]^2 + b_{k+\frac{1}{2}, j-\frac{1}{2}}(I) [\psi_{k+1, j-1} - \psi_{k, j}]^2 \\ &+ b_{k+\frac{3}{2}, j+\frac{1}{2}}(I) [\psi_{k+1, j+1} - \psi_{k+2, j}]^2. \end{aligned} \right.$$

A basic inequality is

Lemma A.4: Let  $\psi \in S_E$  then

$$(A.12a) \quad [\psi_{k+1, j+1} - \psi_{k+1, j-1}]^2 \leq 2\{[\psi_{k+1, j+1} - \psi_{k+2, j}]^2 + [\psi_{k+2, j} - \psi_{k+1, j-1}]^2\}$$

$$(A.12b) \quad [\psi_{k+1, j+1} - \psi_{k+1, j-1}]^2 \leq 2\{[\psi_{k+1, j+1} - \psi_{k, j}]^2 + [\psi_{k, j} - \psi_{k+1, j-1}]^2\}$$

$$(A.12c) \quad [\psi_{k+1, j+1} - \psi_{k+1, j-1}]^2 \leq [\psi_{k+1, j+1} - \psi_{k+2, j}]^2 + [\psi_{k+2, j} - \psi_{k+1, j-1}]^2 \\ + [\psi_{k+1, j+1} - \psi_{k, j}]^2 + [\psi_{k, j} - \psi_{k+1, j-1}]^2$$

$$(A.13a) \quad [\psi_{k+2,j} - \psi_{k,j}]^2 \leq 2\{[\psi_{k+2,j} - \psi_{k+1,j+1}]^2 + [\psi_{k+1,j+1} - \psi_{k,j}]^2\}$$

$$(A.13b) \quad [\psi_{k+2,j} - \psi_{k,j}]^2 \leq 2\{[\psi_{k+2,j} - \psi_{k+1,j-1}]^2 + [\psi_{k+1,j-1} - \psi_{k,j}]^2\}$$

$$(A.13c) \quad [\psi_{k+2,j} - \psi_{k,j}]^2 \leq [\psi_{k+2,j} - \psi_{k+1,j+1}]^2 + [\psi_{k+1,j+1} - \psi_{k,j}]^2 \\ + [\psi_{k+2,j} - \psi_{k+1,j-1}]^2 + [\psi_{k+1,j-1} - \psi_{k,j}]^2 .$$

Proof: Apply the triangle inequality and the inequality  $2ab \leq a^2 + b^2$ .

Lemma A.5: Suppose  $(x_{k+1}, y_j) \in \Omega_h$ . Then there is a constant  $K$  depending only on  $\|\nabla p\|_\infty$  and  $p_0$  such that

$$(A.14) \quad 0 \leq \langle \tilde{L}_E^{(1)} \psi, \psi \rangle_I \leq 2(1+Kh) \langle L_E^{(1)} \psi, \psi \rangle_I .$$

Proof: Note that

$$\frac{1}{2} \langle \tilde{L}_E^{(1)} \psi, \psi \rangle_I = \langle \hat{L}_E \psi, \psi \rangle_I - \frac{1}{2} \langle L_E^{(1)} \psi, \psi \rangle_I .$$

From (A.8c), (A.8d), (A.12a) and (A.12c) we have

$$(A.15a) \quad \frac{1}{2} a_{k+\frac{1}{2}, j+\frac{1}{2}}(I) = \gamma_{k+\frac{1}{2}, j+\frac{1}{2}}(I) ,$$

$$(A.15b) \quad \frac{1}{2} b_{k+\frac{1}{2}, j-\frac{1}{2}}(I) = \sigma_{k+\frac{1}{2}, j-\frac{1}{2}}(I) .$$

From (A.10c), (A.10d), (A.12b), (A.12d) we see that

$$(A.16a) \quad \frac{1}{2} a_{k+\frac{3}{2}, j-\frac{1}{2}}(I) = \gamma_{k+\frac{3}{2}, j-\frac{1}{2}}(I) ,$$

$$(A.16b) \quad \frac{1}{2} b_{k+\frac{1}{2}, j+\frac{1}{2}}(I) = \sigma_{k+\frac{1}{2}, j+\frac{1}{2}}(I) .$$

Therefore from (A.11a) and (A.11b) we have

$$(A.17) \quad \left\{ \begin{aligned} \langle \tilde{L}_E^{(1)} \psi, \psi \rangle_I &= \alpha_{k+1,j}(I) [\psi_{k+2,j} - \psi_{k,j}]^2 \\ &+ \beta_{k+1,j}(I) [\psi_{k+1,j+1} - \psi_{k+1,j-1}]^2. \end{aligned} \right.$$

Thus, we have established the left hand inequality of (A.14). Using (A.12), i.e., the definitions of  $a_{k+\frac{1}{2},j+\frac{1}{2}}(I)$ ,  $b_{k+\frac{1}{2},j-\frac{1}{2}}(I)$  etc. and (A.8b), the definition of  $\alpha_{k+1,j}(I)$  and (A.10b), the definition of  $\beta_{k+1,j}(I)$  we see that there is a constant  $K$ , depending only on  $\|\nabla p\|_\infty$  and  $p_0$  such that

$$(A.18a) \quad \frac{\alpha_{k+1,j}}{a_I} \leq \frac{1}{2}(1+Kh)$$

$$(A.18b) \quad \frac{\beta_{k+1,j}}{a_I} \leq \frac{1}{2}(1+Kh)$$

where

$$(A.18c) \quad a_I = \text{any of } \{a_{k+\frac{1}{2},j+\frac{1}{2}}, a_{k+\frac{3}{2},j-\frac{1}{2}}, b_{k+\frac{1}{2},j-\frac{1}{2}}, b_{k+\frac{3}{2},j+\frac{1}{2}}\}.$$

Therefore (A.17) and (A.12c), (A.13c) and (A.17b) yields the right hand inequality of (A.14).

Corollary:

$$(A.19) \quad \frac{1}{2} \langle L_E^{(1)} \psi, \psi \rangle_I \leq \langle \hat{L}_E \psi, \psi \rangle_I \leq \frac{3}{2}(1+Kh) \langle L_E^{(1)} \psi, \psi \rangle_I.$$

Lemma A.6: Suppose  $(x_{k+1}, y_j) \notin \Omega_h$ . Then the conclusion of lemma A.6 and its corollary hold.

Proof: Calculations similar to those of lemma A.5 now yield

$$(A.20) \quad 0 \leq \langle \tilde{L}_E \psi, \psi \rangle_I = \alpha_{k+1,j} [\psi_{k+2,j} - \psi_{kj}]^2.$$

Thus, as before, the left hand inequality of (A.14) holds. In this case (A.11a), (A.11b) and (A.10c), (A.10d) imply

$$\frac{\alpha_{k+1,j}}{a_I} \leq (1+Kh).$$

Thus, the lemma is proven.

Theorem A: There is a constant  $K$ , depending only on  $\|\nabla p\|_\infty$  and  $p_0$  such that, for both choices of  $L_E (= L_E^{(1)}, L_E^{(2)})$  and associated  $\tilde{L}_E$ , if  $\phi \neq 0$  and  $\phi \in S_E$  we have

$$-Kh \leq \frac{\langle \tilde{L}_E \phi, \phi \rangle}{\langle L_E \phi, \phi \rangle} \leq 2(1+Kh).$$

Proof: The arguments which give lemma A.5 and lemma A.6 extend to all the squares, II, III and IV. Thus, those lemmas imply that the theorem holds for  $L_E = L_E^{(1)}$ . The case of  $L_E = L_E^{(2)}$  follows from (3.5) and the observation that

$$(1-Kh) \langle L_E^{(1)} \psi, \psi \rangle \leq \langle L_E^{(2)} \psi, \psi \rangle \leq (1+Kh) \langle L_E^{(1)} \psi, \psi \rangle.$$

## REFERENCES:

- [1] R. E. Alcouffe, A. Brandt, J. E. Dendy Jr. and J. W. Painter: The Multigrid methods for the diffusion equation with strongly discontinuous coefficients. SIAM J. Sci. Stat. Comput. 2, 430-454 (1981).
- [2] D. Braess: The Contraction Number of a Multigrid Method for Solving the Poisson Equation, Numer. Math., 37, 387-404 (1981).
- [3] A. Brandt: Multilevel adaptive solutions to boundary-value problems, Math. Comp. 31, 333-390 (1977).
- [4] J. E. Dendy, Jr.: Black Box Multigrid, J. Comp. Phys. 48, 366-386 (1982).
- [5] P. O. Frederickson: Fast approximate inversion of large sparse linear systems, Math. Report 7-75, Lakehead University, Ontario, Canada, 1975.
- [6] G. E. Forsythe and W. R. Wasow: Finite Difference Methods for Partial Differential Equations, Wiley Publishers, New York, 1960.
- [7] S. F. McCormick and J. W. Ruge: Multigrid Methods for Variational Problems, SIAM J. Num. Anal., 19, 924-929 (1982).
- [8] M. Ries, U. Trottenberg and G. Winter: A Note on MGR Methods, Linear Algebra Appl. 49, 1-26 (1983).
- [9] J. C. South (Jr.) and A. Brandt: Application of a multi-level grid method to transonic flow calculations. Transonic Flow Problems in Turbomachinery. T. C. Adamson and M. F. Platzer Editors, Hemisphere, Washington D.C.



**END**

**FILMED**

**6-85**

**DTIC**